



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2019

---

## **Alterations in Bone Homeostasis and Microstructure Related to Depression and Allostatic Load**

Wippert, Pia-Maria ; Block, Andrea ; Mansuy, Isabelle M ; Peters, Eva M J ; Rose, Matthias ; Rapp,  
Michael A ; Huppertz, Alexander ; Wuertz-Kozak, Karin

DOI: <https://doi.org/10.1159/000503640>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-176387>

Journal Article

Published Version

Originally published at:

Wippert, Pia-Maria; Block, Andrea; Mansuy, Isabelle M; Peters, Eva M J; Rose, Matthias; Rapp, Michael A; Huppertz, Alexander; Wuertz-Kozak, Karin (2019). Alterations in Bone Homeostasis and Microstructure Related to Depression and Allostatic Load. *Psychotherapy and Psychosomatics*, 88(6):383-385.

DOI: <https://doi.org/10.1159/000503640>

Psychother Psychosom  
DOI: 10.1159/000503640

## Alterations in Bone Homeostasis and Microstructure Related to Depression and Allostatic Load

Pia-Maria Wippert<sup>a,b</sup> Andrea Block<sup>a</sup> Isabelle M. Mansuy<sup>c</sup>  
Eva M.J. Peters<sup>d,e</sup> Matthias Rose<sup>f</sup> Michael A. Rapp<sup>g</sup>  
Alexander Huppertz<sup>h</sup> Karin Wuertz-Kozak<sup>a,b,i-k</sup>

<sup>a</sup>Sociology of Health and Physical Activity, Department of Health Sciences, University of Potsdam, Potsdam, Germany; <sup>b</sup>Institute for Biomechanics, Department of Health Sciences and Technology, ETH Zurich, Zurich, Switzerland; <sup>c</sup>Laboratory of Neuroepigenetics, University of Zurich/Swiss Federal Institute of Technology, Brain Research Institute, Neuroscience Center Zurich, Zurich, Switzerland; <sup>d</sup>Psychoneuroimmunology Laboratory, Department of Psychosomatics and Psychotherapy, University of Giessen, Giessen, Germany; <sup>e</sup>Charité Center 12, Internal Medicine and Dermatology, Division for General Internal Medicine, Psychosomatics and Psychotherapy, Universitätsmedizin Berlin, Berlin, Germany; <sup>f</sup>Charité, Division for General Internal Medicine, Psychosomatics and Psychotherapy, Universitätsmedizin Berlin, Berlin, Germany; <sup>g</sup>Social and Preventive Medicine, Department of Health Sciences, University of Potsdam, Potsdam, Germany; <sup>h</sup>University of Potsdam, Outpatient Clinic, Center of Sports Medicine, Potsdam, Germany; <sup>i</sup>Schön Clinic Munich Harlaching, Spine Center and Academic Teaching Hospital of the Paracelsus Medical University, Salzburg, Austria; <sup>j</sup>Schön Clinic Munich Harlaching, Spine Center and Academic Teaching Hospital of the Paracelsus Medical University, Munich, Germany; <sup>k</sup>Department of Biomedical Engineering, Rochester Institute of Technology (RIT), Rochester, NY, USA

Depression is a severe and highly comorbid condition that affects mental and physical health and is a prime cause of disability. Regarding the pathogenesis of comorbid depressive disorders, previous research has shown that neuroendocrine and neuroanatomical responses to stress lead to cellular and molecular changes that promote the development of comorbid diseases [1]. Hence, depressive disorders may influence the homeostasis of tissues, including bone, and contribute, for example, to osteoporosis development [2, 3] and decreased bone mineral density (BMD) [4]. However, the present knowledge of involved pathogenetic mechanisms is still scarce, and studies are required that provide a better understanding of possible interactions between depression, neuroendocrine stress responses, and bone metabolism. Moreover, it is unclear what comes first: does mental distress, such as depres-

sion, impair bone health or does stress-induced disruption of bone homeostasis come first? Both perspectives are highly relevant for the medical care of patients [4], and their study requires the definition of relevant outcomes.

Present stress concepts state that neuroendocrine signaling pathways are activated by stress in an attempt to enable mental as well as physical adaptation and maintain a low allostatic load. However, persistent excessive or prolonged stresses (e.g., traumatic life events, chronic work strain triggering a depressive episode) can provoke a maladaptive increase of neuroendocrine responses, toxic systemic stress, and an accumulation of allostatic load [5]. It is evident that specific allostatic load mediators, such as cortisol, growth hormone, or insulin-like growth factor-I, may play a role in the lifelong remodeling process of bones [6, 7]. Elevated cortisol levels can inhibit osteoblast proliferation, differentiation, and apoptosis, leading to decreased bone density. However, whether such allostatic overload occurs depends largely on exposure intensity and duration, personal and social resources (coping, exposure), and epigenetic adaptability [8]. In other words, stressful life events do not necessarily lead to depressive disorders or allostatic overload, especially if the neuroendocrine response enables adaptation and can be turned off after an appropriate interval [9].

Here, we therefore aimed to study whether bone metabolism is changed during an acute depressive episode and whether people with a high life burden and allostatic load show alterations in bone metabolism and mineral density. Further, we wanted to know whether a mathematical algorithm can determine individual allostatic load mediators that identify patients at higher risk for disrupted bone metabolism and thereby optimize medical care.

In an 18-month, multicenter, observational study, we assembled a cohort of two hundred and eight 18- to 65-year-old depressive patients in the primary care setting suffering from severe to moderate depression (mainly ICD-10 F32.x or F33.x). We determined psychosomatic symptoms (Beck Depression Inventory-II [BDI-II]; Symptom Check List – revised [SCL-90-R]; Positive Symptom Distress Index [PSDI], Global Severity Index [GSI], and Positive Symptom Total [PST]), life burden (Inventory of Life Changing Events), and allostatic load index (ALI; derived from 15 neuroendocrinological, immunological, and metabolic allostatic mediators). Bone metabolism was measured by 3 bone formation and resorption serum markers (osteocalcin [OC], procollagen type 1 N-terminal propeptide [P1NP], and  $\beta$ -CrossLaps [CTX]). BMD was assessed in both hips and the lumbar spine (subsample,  $n = 18$ ; DXA BMD-L1L4, -TMW, -SMW, -GMW, and -HMW). Descriptive statistics, paired sample  $t$  tests, multivariable regression models, and Least Absolute Shrinkage and Selection Operator (LASSO) were applied for analyzing the study questions.

65% of our patients (mean age = 46.63, SD = 11.13,  $f = 75\%$ ) suffered from severe to moderate depression (mean BDI-II = 23.61, SD = 10.14), severe psychosomatic symptoms (GSI: 41.8%; PSDI: 42.3%), and allostatic load burden (ALI: 29.3%). Higher life

**Table 1.** Main effects (regression coefficient *b*) of depressiveness, life events, psychosomatic symptoms severity, and ALIs on bone markers at baseline (*t*<sub>0</sub>)

	P1NP			OC			CTx		
	coefficient	95% CI	<i>p</i>	coefficient	95% CI	<i>p</i>	coefficient	95% CI	<i>p</i>
BDI-II sum score	0.121	−0.170, 0.413	0.417	<b>0.083</b>	−0.008, 0.173	0.075	0.0004	−0.002, 0.002	0.727
SCL-90-R PSDI	<b>5.304</b>	−0.585, 11.194	0.081	<b>2.631</b>	0.855, 4.407	0.005	<b>0.052</b>	0.011, 0.093	0.016
SCL-90-R PST	0.061	−0.134, 0.256	0.543	0.045	−0.015, 0.105	0.145	−0.0002	−0.002, 0.001	0.729
SCL-90-R GSI	3.222	−2.103, 8.548	0.239	<b>2.096</b>	0.447, 3.745	0.015	0.020	−0.020, 0.059	0.333
ILE all <sup>a</sup>	0.158	−0.153, 0.470	0.317	<b>0.111</b>	0.017, 0.206	0.021	<b>0.002</b>	0.000, 0.004	0.049
ALI 21	−0.523	−1.650, 0.604	0.365	− <b>0.438</b>	−0.779, −0.096	0.014	− <b>0.009</b>	−0.017, −0.001	0.025
ALI 15	−0.435	−1.759, 0.888	0.521	− <b>0.462</b>	−0.864, −0.059	0.027	− <b>0.009</b>	−0.018, 0.0005	0.065
ALI-II	−0.351	−1.585, 0.884	0.579	− <b>0.492</b>	−0.875, −0.108	0.014	− <b>0.008</b>	−0.017, 0.001	0.088
ALI-I	0.734	−1.824, 3.291	0.576	0.467	−0.318, 1.253	0.246	−0.005	−0.023, 0.014	0.614

Multiple interaction regression models, adjusted for age, gender, and study sites. Significant regression coefficients are bold ( $p < 0.01$ ,  $p < 0.05$ ,  $p < 0.10$ ; 2-sided testing). ALI, allostatic load index; BDI-II, Beck Depression Inventory II; SCL-90-R, Symptom Check List – revised; PSDI, Positive Symptom Distress Index; PST, Positive Symptom Total; GSI, Global Severity Index; ILE, Inventory of Life Changing Events. <sup>a</sup> Count of all life events.

burden was associated with depressive disorder (BDI-II:  $b = 0.24$  [95% CI 0.064, 0.411],  $p = 0.008$ ) but not with allostatic load.

In this cohort, we observed that higher depression (OC:  $b = 0.08$  [95% CI −0.008, 0.173],  $p = 0.075$ ) and symptom severity (PSDI: P1NP:  $b = 5.30$  [95% CI −0.59, 11.19],  $p = 0.081$ ; OC:  $b = 0.91$  [95% CI 0.86, 4.41],  $p = 0.005$ ; CTx:  $b = 0.052$  [95% CI 0.011, 0.093],  $p = 0.016$ ; GSI: OC:  $b = 2.10$  [95% CI 0.45, 3.75],  $p = 0.015$ ) were associated with a higher expression of anabolic bone metabolism (OC; P1NP; CTx). People with a high life burden showed significantly less anabolic expression (OC:  $b = 0.11$  [95% CI 0.017, 0.206],  $p = 0.021$ ; CTx:  $b = 0.002$  [95% CI 0.000, 0.004],  $p = 0.049$ ).

At the same time, patients with high allostatic load showed a significantly decreased bone metabolism (ALI: OC:  $b = −0.46$  [95% CI −0.864, −0.059],  $p = 0.027$ ; CTx:  $b = −0.01$  [95% CI −0.018, 0.001],  $p = 0.065$ ) and reduced BMD in the lumbar spine and both hips (DXA). LASSO could not identify any singular allostatic load mediator sufficient enough for the prediction of bone metabolism quality (OC; P1NP; CTx) 1.5 years after baseline (Table 1).

The findings suggest that anabolic activation of bone metabolism occurs during a depressive episode, which increases with the severity of the depression and psychosomatic symptoms. This so far undiscovered metabolic adaptation to an exceptionally high stress episode is limited to patients with a higher life burden and completely absent in patients with allostatic overload. The latter are – presumably due to the accumulation of toxic stress in tissues and cells – no longer capable of an adequate metabolic adaptation. Thus, a catabolic and bone damaging process occurs, which is reflected by a reduction of BMD in the long term.

The fact that predominantly patients with acute and nonchronic depressive disorder were involved in the study enabled the discovery of this highly important mechanism that is believed to affect corresponding medical therapies (drug administration, exercise, etc.). The data provide further evidence that a gradation between different levels of physical and mental load accumulation must be respected. However, many questions remain unanswered and should be investigated in further studies. For example, no diagnos-

tic instrument could be derived from the allostatic mediators. In addition, further studies should examine to what extent psychometric data would allow diagnostic assessments [9].

#### Acknowledgement

We are grateful to our statistician David Drießlein for his independent analysis of the data and to Sanne Houtenbos for proof reading. Furthermore, we thank our colleagues from the DEPRE-HA network, i.e., Andreas Häusler, Friederike Deeken, Susanne Schulze, Michael Rector, and Christine Wiebking. We thank the technical and medical staff of the study sites for their contributions throughout the study.

#### Statement of Ethics

All clinical investigations were conducted according to the principles expressed in the Declaration of Helsinki. Final ethical approval was provided on December 11, 2017, from the major institutional ethics review board of the University of Potsdam, Germany (number 15/2017). All participants gave their written informed consent.

#### Disclosure Statement

The authors have no conflicts of interest to declare.

#### Funding Sources

The present study was funded by the University of Potsdam's "Central Research Grant for Innovative Projects." We further acknowledge the support of the Deutsche Forschungsgemeinschaft, the Open Access Publishing Fund of the University of Potsdam, and the German Pension Fund Association (10-R-40.07.05.07.008).

#### Author Contributions

P.-M.W., K.W.-K.: conceptualization and first draft; P.M.-W., K.W.-K., A.B.: data analysis and interpretation; P.M.-W., A.B., M.R., A.H.: data acquisition. P.M.-W., A.B., I.M.M., E.M.J.P., M.R., A.H., M.A.R., and K.W.-K. substantially improved and revised the manuscript. All authors approved the final version.

## References

- 1 Gold PW. The organization of the stress system and its dysregulation in depressive illness. *Mol Psychiatry*. 2015 Feb;20(1):32–47.
- 2 Picard M, McEwen BS, Epel ES, Sandi C. An energetic view of stress: focus on mitochondria. *Front Neuroendocrinol*. 2018 Apr;49:72–85.
- 3 Wippert PM, Rector M, Kuhn G, Wuertz-Kozak K. Stress and Alterations in Bones: An Interdisciplinary Perspective. *Front Endocrinol (Lausanne)*. 2017 May;8(96):96.
- 4 Schweiger JU, Schweiger U, Hüppe M, Kahl KG, Greggersen W, Fassbinder E. Bone density and depressive disorder: a meta-analysis. *Brain Behav*. 2016 May;6(8):e00489.
- 5 McEwen BS. Allostasis and the epigenetics of brain and body health over the life course: the brain on stress. *JAMA Psychiatry*. 2017 Jun;74(6):551–552.
- 6 Zhang M, Xuan S, Bouxsein ML, von Stechow D, Akeno N, Faugere MC, et al. Osteoblast-specific knockout of the insulin-like growth factor (IGF) receptor gene reveals an essential role of IGF signaling in bone matrix mineralization. *J Biol Chem*. 2002 Nov;277(46):44005–12.
- 7 Furlan PM, Ten Have T, Cary M, Zemel B, Wehrli F, Katz IR, et al. The role of stress-induced cortisol in the relationship between depression and decreased bone mineral density. *Biol Psychiatry*. 2005 Apr;57(8):911–7.
- 8 McEwen BS, Nasca C, Gray JD. Stress Effects on Neuronal Structure: Hippocampus, Amygdala, and Prefrontal Cortex. *Neuropsychopharmacology*. 2016 Jan;41(1):3–23.
- 9 Fava GA, McEwen BS, Guidi J, Gostoli S, Offidani E, Sonino N. Clinical characterization of allostatic overload. *Psychoneuroendocrinology*. 2019 Oct;108:94–101.